

Reverse Mutations in the Fragile X Syndrome

W. Ted Brown, George E. Houck, Jr., Xiaohua Ding, Nan Zhong, Sarah Nolin, Anne Glicksman, Carl Dobkin, and Edmund C. Jenkins

New York State Institute for Basic Research in Developmental Disabilities, Staten Island, New York

Three females were identified who have apparent reversal of fragile X premutations. Based on haplotype analysis of nearby markers, they were found to have inherited a fragile X chromosome from their premutation carrier mothers, and yet had normal size *FMR1* repeat alleles. The changes in repeat sizes from mother to daughter was 95 to 35 in the first, 145 to 43 in the second, and 82 to 33 in the third. In the first family, mutations of the nearby microsatellites *FRAXAC2* and *DXS548* were also observed. In the other two, only mutations involving the *FMR1* repeats were found. We suggest differing mutational mechanisms such as gene conversion versus DNA replication slippage may underlie such reversions. We estimate that such revertants may occur among 1% or less of premutation carrier offspring. Our results indicate that women identified to be carriers by linkage should be retested by direct DNA analysis.

© 1996 Wiley-Liss, Inc.

KEY WORDS: fragile X syndrome, DNA repetitive sequences, mutation

INTRODUCTION

The fragile X syndrome is the most common inherited form of mental retardation [Brown and Jenkins, 1992; Warren and Nelson, 1994]. Previously, the diagnosis of the syndrome was established by the cytogenetic demonstration of a fragile site at Xq27.3 (*FRAXA*) under conditions of folate deprivation. The cloning and characterization of the fragile X gene (*FMR1*) led to direct DNA testing and recognition that the mutation responsible for the syndrome generally involves expansion of a repeated trinucleotide (CGG)_n sequence

located within the gene's 5' non-coding region [Verkerk et al., 1991; Fu et al., 1991; Rousseau et al., 1991]. The normal number of repeats varies from approximately 6 to 55 with the most common Caucasian allele size being 30 [Brown et al., 1993; Snow et al., 1993]. Normal male and female carriers of the mutation generally have from approximately 56 to 200 repeats [Fu et al., 1991; Brown et al., 1993; Snow et al., 1993]. Such individuals are now commonly referred to as carriers of a "premutation." The concept of a premutation with regard to fragile X was initially suggested by Sherman et al. [1985] and by Pembrey et al. [1985] in the context of attempting to model the apparent delayed mutations seen in fragile X families. Premutations are generally considered to have a risk to expand to the full mutation in the next generation or in the second generation in the case of a transmitting male and are usually ascertained in the context of a family study of a full mutation fragile X index case [Fu et al., 1991; Warren and Nelson, 1994]. Full mutations as found in affected individuals generally have more than ~200 repeats are usually associated with methylation of an upstream promoter region and have a lack of gene expression [Pierretti et al., 1991]. The risk of expansion from a premutation to a full mutation is dependent upon sex of the parent, the size of the repeat, and by lack of interruptions of the pure CGG repeat with AGGs that may increase stability [Eichler et al., 1994; Kunst and Warren, 1994; Snow et al., 1994; Hirst et al., 1994; Zhong et al., 1995]. No new fragile X mutations have been described [Smits et al., 1993]. Because of this low frequency of new mutations, the fragile X syndrome is atypical of other severe X-linked diseases with reduced genetic fitness where new mutation rates are generally high to replace the loss of disease alleles.

Previously, we noted a family in which a reverse fragile X mutation was apparent in a daughter [Zhong et al., 1993]. We have now identified two more such females. These three unrelated females inherited the same flanking fragile X chromosome markers as their affected male brothers. In each case they were initially given high risks of being carriers, based on linkage analysis. However, direct mutation testing has shown each to have two normal-sized *FMR1* alleles. Thus, these three women have alleles that represent apparent reverse fragile X mutations.

Received for publication September 19, 1995; revision received December 21, 1995.

Address reprint requests to Dr. W. Ted Brown, Department of Human Genetics, Institute for Basic Research in Developmental Disabilities, 1050 Forest Hill Road, Staten Island, NY 10314.

MATERIALS AND METHODS

Blood specimens were collected in EDTA tubes and DNA was extracted by phenol/chloroform or by using an extraction kit (Gentra, Inc). Testing for flanking RFLPs was done using probes and of microsatellite repeats by PCR using the referenced procedures. The closely flanking polymorphic loci that were typed included *DXS539* (pJH89) [van den Hurk et al., 1991], *DXS297* (VK23) [Suthers et al., 1991], *DXS548* [Riggins et al., 1992], *FRAXAC1* [Richards et al., 1991], *FMRA* [Kunst and Warren, 1994], *FRAXAC2* [Richards et al., 1991], *DXS465* (Do33) [Oberlé et al., 1991], *FRAXE* [Knight et al., 1993], *DXS296* (VK21) [Suthers et al., 1991], and *DXS374* (1A1) [Patterson et al., 1989]. The locations of the polymorphic loci used for haplotyping are indicated in Figure 1. Direct Southern blot analysis of the *FMR1* locus following digestions with enzymes *EcoRI* and *EagI* used probe StB12.3 following the method of Rousseau et al. [1991]. *FMR1* CGG repeat numbers were determined using our PCR method employing forward primer 1 and reverse primer 3 as previously described [Brown et al., 1993]. Cytogenetic procedures for fragile X detection were performed as previously described [Jenkins et al., 1992].




Fig. 1. Approximate location of DNA markers used relative to the *FMR1* gene and CGG repeat on the X chromosome.




Fig. 2. Family 1 with DNA haplotypes. For Figures 2–4, the fragile X chromosomes are indicated in black, the maternal non-fragile X in gray and the paternal chromosome in white. The location of the reverse mutations of *FMR1* CGG repeats and of microsatellite mutations are indicated by stars.

RESULTS

The three family pedigrees and the informative markers studied are illustrated in Figures 2–4. *FMR1* allele PCR analysis is shown in Figure 5. Southern blot analysis using probe StB12.3 confirmed that the affected males had typical full mutations, that their mothers were premutation carriers, and that none of the three daughters showed a premutation pattern. Cytogenetic testing confirmed that at least one affected male in each family was positive for the fragile X chromosome.




Fig. 3. Family 2 with DNA haplotypes.




Fig. 4. Family 3 with DNA haplotypes.

Family 1

The first family included the carrier mother, I-1, with *FMR1* alleles of 30 and 95 repeats, the father, I-2, with 20 repeats, one affected son, II-1, with a full mutation,

one normal daughter, II-2, with 20 and 35 repeats, and one normal son, II-3, with 30 repeats (Fig. 2). The daughter had three normal children (not shown), all of whom had inherited her 20 repeat allele. The daughter had inherited the same set of flanking markers from her carrier mother as that of her affected brother, including *FRAXAC1* and *DXS465*. Thus, she had inherited the fragile X region from her carrier mother and the mother's premutation allele had decreased from 95 to 35 repeats in the daughter. Previously, we briefly noted this family and that mutations were also observed in *FRAXAC2* [Zhong et al., 1993]. The carrier mother had *FRAXAC2* allele lengths of 146 and 149 bp. Her affected and her normal sons both had 150 bp alleles, while the daughter had inherited a 147 bp allele [Zhong et al., 1993]. Sequencing of *FRAXAC2* for the two brothers showed both had the same allele (X-16, Y-7, Z-14). In addition, the normal brother also had a 194 bp *DXS548* allele while the mother was homozygous for 196 indicating three loci showed mutations in this family. Repeat samplings on this family confirmed the typings were correct.

Family 2

The second family included a carrier mother, I-1, with *FMR1* alleles of 29 and 145 repeats, a father, I-2, with 30 repeats, one affected son, II-1, with a full mutation, one daughter, II-2, with 30 and 43 repeats, and a second normal daughter, II-3, with 29 and 30 repeats




Fig. 5. PCR analysis of repeat sizes for the three families showing reverse mutations for the daughters. Pedigree numbers and lanes correspond to Figures 2-4. The daughters with reverse mutations are each indicated as II-2. All samples were analyzed on the same autoradiogram with a 1 hour exposure. Overnight exposure was required to better visualize the marker lanes (M) and some alleles (family 2: I-1, I-2; family 3, I-1).

(Fig. 3). Based on six informative flanking markers, the daughter, II-2, was found to have inherited the same fragile X chromosome as her affected brother indicating the mother's 145 repeat allele had undergone a reduction to the 43 repeat allele during transmission.

Family 3

The third family included a carrier mother, I-1, with *FMR1* alleles of 30 and 82 repeats, a father, I-2, with 23 repeats, one affected son, II-1, with a full mutation, one daughter, II-2, with 23 and 33 repeats, and a second daughter, II-3, with 23 and 30 repeats (Fig. 4). The carrier mother was informative for the two closest flanking markers, *FRAXAC1* and *FMRa*, as well as *DXS548* and *FRAXAC2*. These markers indicated that daughter II-2 had inherited the same fragile X chromosome as her affected brother but that she had a 33 repeat allele instead of an 82 repeat allele or a larger size indicating a reduction in length had occurred. We obtained a buccal epithelial sample for this daughter and typing confirmed she had allele sizes of 23 and 33.

DISCUSSION

These three examples of apparent reverse mutations are the only we have identified having analyzed the offspring of approximately 300 fragile X female premutation carriers. These events are not common and we estimate they may occur among 1% or less of meioses. These three women have normal-size but potentially unstable alleles and prenatal diagnosis may be advisable for their future pregnancies. Our results suggest that women who were previously counseled to have a high risk of being a carrier based on linkage analysis should be retested with *FMR1* mutational analysis to confirm their status.

Detection of such reverse mutations would be likely only if previous linkage studies and direct testing were discrepant or if PCR sizing analysis revealed discrepancies between parents and child. Since none of these daughters in these families has yet passed the altered repeat to the next generation, the stability of these alleles is yet unknown. The buccal sample in the one case tested was identical to her blood sample suggesting no somatic mosaicism was present. While mosaicism in which a premutation allele is present in the germline can not yet be disproven, we believe the risk for these women to bear a child with the full mutation is likely to be greatly reduced, since no full mutation child has been identified having a mother with an allele size of <55 repeats (J. Holden, these proceedings). We have found that approximately two thirds of premutation alleles lack any interrupting AGGs and the remainder have only one, and thus these three women are most likely to have a pure CGG repeats of 33, 35, and 43 [Zhong et al., 1995]. We lack confirmation of this conclusion, since we have not yet been able to adopt our method for AGG analysis to females. Further, cloning and subsequent sequencing, although possible, may be unreliable because of rearrangement artifacts. Instabilities of allele transmissions within the range of 36 to

55 repeats have been described indicating there exists a "gray zone" of potentially unstable alleles [Eichler et al., 1994, Zhong et al., 1995, 1996 these proceedings]. Which gray zone alleles should be considered as "pre-mutations" in the absence of an identified close relative with a full mutation is unclear.

These events did not appear to involve a deletion of regions flanking the repeat, since the mutated alleles could be amplified by PCR primers that closely flank the CGG repeat region (ending 16 bp upstream and 5 bp downstream). Rather, the mutational change in each case appeared to be within the CGG repeat. In family one, both the CGG repeat and the *FRAXAC2* locus underwent mutational changes in two siblings. Since these loci are only separated by ~12 kb, the occurrence of related single events, such as gene conversion, seems likely. (The only known intervening polymorphism, *FMRa*, was uninformative, and thus we can not determine whether an unequal recombination event or whether two independent mutations occurred). Two microsatellite mutations involving *DXS548* and *FRAXAC2* were also observed in the normal male sibling in this family, but the *FMR1* allele remained unchanged, suggesting two double recombinations or a complex gene conversion. The reverse mutations in the second and third cases appeared to be confined to a change within the CGG repeat, as both of the closest upstream and downstream polymorphisms, *FRAXAC1* and *FMRa*, were the same as their affected brothers. Since there was no apparent recombination or mutation of flanking markers, the latter could be due to a simpler mechanism such as DNA slippage.

Three other instances of apparent fragile X reverse mutations have been reported. In the first [Snow et al., 1993], a daughter who had been given a high risk to be a carrier by linkage was found by Southern analysis to have two normal size alleles. In the second [Vits et al., 1994], a carrier mother with a 110 repeat allele transmitted a 44 repeat allele to her daughter. In the third [van den Ouweland et al., 1994], a carrier mother with alleles of 80 and 29 repeats transmitted a 29 repeat allele to her daughter. In the latter, proximal and distal markers with one exception (*FMRb*) were the same as the affected haplotype and a complex mutation involving multiple exchanges was suggested. Since each of the six identified revertants have occurred in daughters and no such revertants have been described yet among sons, a sex-related difference in mutational mechanisms may exist.

Gene conversion events appear to be very common at some human minisatellite loci, such as at MS32 [Jeffreys et al., 1994], in which both simple conversions as well as more complex multistep mutation processes involving allele reduplication plus inter-allelic conversion events have been identified. Gene conversion events between similar sequences may also be common, as is suggested by a high frequency of conversions between unlinked, hemizygous transgenes in the germline of mice [Murti et al., 1994]. Similar to fragile X, there have been several examples of reverse mutations described for myotonic dystrophy (DM). Analysis of one indicated a complex gene conversion event

[O'Hoy et al., 1993]. In two others the flanking markers were unchanged [Brunner et al., 1994], suggesting slippage.

There appears to be a size threshold of around 40 pure CGG triplet repeats above which *FMR1* allele instability is much more likely. As we first suggested [Brown et al., 1993], this threshold may correspond to the upper size limit of Okazaki fragments which could increase the probability of DNA slippage. This hypothesis has been further discussed [Kunst and Warren, 1994; Eichler et al., 1994; Richards and Sutherland, 1994]. It has also been noted that 40 repeats is the approximate threshold for the formation of stable hairpin loops which could promote expansion [Chen et al., 1995; Gacy et al., 1995]. Complex internal mechanisms that involve multistep processes resulting in rearrangement of flanking markers also could underlie such mutations [Jeffreys et al., 1994]. Analysis of additional such revertants may shed light on the molecular mechanisms which underlie triplet repeat instability.

ACKNOWLEDGMENTS

We are indebted to Ponmani Goonewardena, Allyn McConkie-Rosell, Kathleen W. Rao, Barbara Wedehase, and Celeste Krauss for assistance with the family studies and Stephanie Sherman for helpful discussions. This study was supported by the New York State Office of Mental Retardation and Developmental Disabilities and by grants from the MCH (MCJ360587) and NIH (HD29407).

REFERENCES

- Brown WT, Jenkins EC (1992): The fragile X syndrome. In T Friedmann (ed): "Molecular Genetic Medicine." New York: Academic Press, Inc., Vol II, pp 39-66.
- Brown WT, Houck G Jr, Jeziorowska A, Levinson F, Ding X-H, Dobkin C, Zhong N, Henderson J, Sklower-Brooks S, Jenkins EC (1993): Rapid fragile X carrier screening and prenatal diagnosis by a non-radioactive PCR test. *JAMA* 270:1569-1575.
- Brunner HG, Jansen G, Nillesen W, Nelen MR, de Die CEM, Höweler CJ, van Oost BA, Wieringa B, Ropers H-H, Smeets HJM (1994): Reverse mutation in myotonic dystrophy. *N Engl J Med* 328:476-480.
- Chen X, Mariappan VS, Castasti P, Ratliff R, Moyzis RK, Laayoun A, Smith SS, Bradbury EM, Gupta G (1995): Hairpins are formed by the single DNA strands of the fragile X triplet repeats: Structure and biological implications. *Proc Natl Acad Sci USA* 92:5199-5203.
- Eichler EE, Holden JJA, Popovich BW, Reiss AL, Snow K, Thibodeau SN, Richards CS, Ward PA, Nelson DL (1994): Length of uninterrupted CGG repeats determines instability in the *FMR1* gene. *Nat Genet* 8:88-94.
- Fu YH, Kuhl DPA, Pizzuti A, Pieretti M, Sutcliffe JS, Richards S, Verkerk A, Holden JJA, Fenwick RG Jr., Warren ST, Oostra BA, Nelson DL, Caskey CT (1991): Variation of the CGG repeat at the fragile X site results in genetic instability: Resolution of the Sherman paradox. *Cell* 67:1047-1058.
- Gacy AM, Goellner G, Juranic N, Macura S, McMurray CT (1995): Trinucleotide repeats that expand in human disease form hairpin structure in vitro. *Cell* 81:533-540.
- Hirst MC, Grewal PK, Davies KE (1994): Precursor arrays for triplet repeat expansion at the fragile X locus. *Hum Mol Genet* 3:1553-1560.
- Jeffreys AJ, Tamaki K, MacLeod A, Monckton DG, Neil DL, Armour JAL (1994): Complex gene conversion events in germline mutation at human minisatellites. *Nat Genet* 6:136-145.
- Jenkins EC, Brown WT, Schonberg S, Krawczun MS, Goldberg J, Golbus MS (1992): Prenatal detection of fra(X)(q27.3) in female identical twins: Reliability of low level cytogenetic prenatal expression in females. *Am J Med Genet* 43:128-35.
- Knight SJL, Flannery AV, Hirst MC, Campbell, Christodoulou Z, Phelps SR, Pointon J, Middleton-Price HR, Barnicoat A, Pembrey ME, Holland J, Oostra BA, Bobrow M, Davis KE (1993): Trinucleotide repeat amplification and hypermethylation of a CpG island in FRAXE mental retardation. *Cell* 74:127-134.
- Kunst CB, Warren ST (1994): Cryptic and polar variation of the fragile X repeat could result in predisposing normal alleles. *Cell* 77:853-861.
- Murti JR, Bumbulis M, Schiementi JC (1994): Gene conversion between unlinked sequences in the germline of mice. *Genetics* 137:837-843.
- Oberlé I, Rousseau F, Heitz D, Kretz C, Devys D, Hanauer A, Boué J, Bertheas MF, Mandel J-L (1991): Instability of a 550-base pair DNA segment and abnormal methylation in fragile X syndrome. *Science* 252:1097-1102.
- O'Hoy KL, Tsilfidis C, Mahadevan M, Neville CE, Barceló J, Hunter AGW, Korneluk RG (1993): Reduction in size of the myotonic dystrophy trinucleotide repeat mutation during transmission. *Science* 259:809-12.
- Patterson MN, Bell MV, Bloomfield J, Flint T, Dorkins H, Thibodeau S, Schaid D, Bren G, Schwartz C, Wieringa B, Ropers H, Callen D, Sutherland G, Froster-Iskenius U, Vissing H, Davies K (1989): Genetic and physical mapping of a novel region close to the fragile X site on the human X chromosome. *Genomics* 4:570-578.
- Pembrey ME, Winter RM, Davies KE (1985): A premutation that generates a defect at crossing over explains the inheritance of fragile X mental retardation. *Am J Med Genet* 21:709-717.
- Pieretti M, Zhang F, Fu YH, Warren ST, Oostra BA, Caskey CT (1991): Absence of expression of the *FMR-1* Gene in fragile X syndrome. *Cell* 66:1-20.
- Richards RI, Holman K, Kozman H, Kremer E, Lynch M, Pritchard M, Yu S, Mulley J, Sutherland GR (1991): Fragile X syndrome: Genetic localization by linkage mapping of two microsatellite repeats FRAXAC1 and FRAXAC2 which immediately flank the fragile site. *J Med Genet* 28:818-23.
- Richards RI, Sutherland GR (1994) Simple repeat DNA is not replicated simply. *Nat Genet* 6:114-116.
- Riggins GJ, Sherman SL, Oostra BA, Sutcliffe JS, Feitell D, Nelson DL, van Oost BA (1992): Characterization of a highly polymorphic dinucleotide repeat 150 kb proximal to the fragile X site. *Am J Med Genet* 43:237-243.
- Rousseau F, Heitz D, Biancalana V, Blumenfeld S, Kretz C, Boué J, Tommerup N, Van Der Hagen C, DeLozier-Blanchet C, Croquette M-F, Gilgenkrantz S, Jalbert P, Voelckel M-A, Oberlé I, Mandel J-L (1991): Direct diagnosis by DNA analysis of the fragile X syndrome of mental retardation. *N Engl J Med* 325:1673-1681.
- Sherman SL, Jacobs PA, Morton NE, Faaroster-Iskenius U, Howard-Peebles PN, Nielsen KB, Partington MW, Sutherland GR, Turner G, Wilson M (1985): Further segregation analysis of the fragile X syndrome with special reference to transmitting males. *Hum Genet* 69:289-299.
- Smits APT, Dreesen JCFM, Post JG, Smeets DFCM, de Die-Smulders C, Spaans-van der Bijl T, Govaerts LCP, Warren ST, Oostra B, van Oost BA (1993): The fragile X syndrome: no evidence for any recent mutations. *J Med Genet* 30:94-96.
- Snow K, Doud LK, Hagerman R, Pergolizzi RG, Erster SH, Thibodeau SN (1993): Analysis of a CGG sequence at the *FMR-1* locus in fragile X families and in the general population. *Am J Hum Genet* 53:1217-28.
- Snow K, Tester DJ, Kruckeberg KE, Schaid DJ, Thibodeau SN (1994): Sequence analysis of the fragile X trinucleotide repeat: Implications for the origin of the fragile X mutation. *Hum Mol Genet* 3:1543-1551.
- Suthers GK, Mulley JC, Voelckel MA, Dahl N, Väisänen ML, Steinback P, Glass IA, Schwartz CE, van Oost BA, Thibodeau SN, Haites NE, Oostra BA, Giné R, Carballo M, Morris CP, Hopwood JJ, Sutherland GR (1991): Genetic mapping of new DNA probes at Xq27 defines a strategy for DNA studies in the fragile X syndrome. *Am J Hum Genet* 48:460-467.
- van den Hurk JAJM, Dreesen JCFM, van den Berg H, van Bennekou CA, van den ouweland AMW, van Oost BA (1991): Two polymorphisms at the DXS539 locus. *Nucl Acid Res* 19:1723.

- van den Ouweland AMW, Deelen WH, Kunst CB, Uzielli ML G, Nelson DL, Warren ST, Oostra BA, Halley DJJ (1994): Loss of mutation at the *FMR1* locus through multiple exchanges between maternal X chromosomes. *Hum Mol Genet* 3:1823-1827.
- Verkerk AJMH, Pieretti M, Sutcliffe JS, Fu YH, Kuhl DPA, Pizzuti A, Reiner O, Richards S, Victoria MF, Zhang F, Eussen BE, van Ommen GJB, Blonden LAJ, Riggins GJ, Chastain JL, Kunst CB, Galjaard H, Caskey CT, Nelson DL, Oostra BA, Warren ST (1991): Identification of a gene (*FMR-1*) containing a CGG repeat coincident with a breakpoint cluster region exhibiting length variation in fragile X syndrome. *Cell* 65:905-14.
- Vits L, De Boulle K, Reyniers E, Handig I, Darby JK, Oostra B, Willems PJ (1994). Apparent regression of the CGG repeat in *FMR1* to an allele of normal size. *Hum Genet* 94:523-526.
- Warren ST, Nelson DL (1994): Advances in molecular analysis of fragile X syndrome. *JAMA* 271:536-542.
- Zhong N, Dobkin C, Brown WT (1993): A complex mutable polymorphism located within the fragile X gene. *Nat Genet* 5:248-253.
- Zhong N, Yang WH, Dobkin C, Brown WT (1995) Fragile X gene instability: Anchoring AGGs and linked microsatellites. *Am J Hum Genet* 57:351-361.
- Zhong N, Ju W, Pietrofesa J, Wang D, Dobkin C, Brown WT (1996): Fragile X "gray zone" alleles: AGG patterns, expansion risks and associated haplotypes. *Am J Med Genet* 64:261-265.